






Metagenome-Assembled Genome Sequences Obtained from a Reactivated Kombucha Microbial Community Exposed to a Mars-Like Environment outside the International Space Station

Imchang Lee,^{a,b} Debmalya Barh,^c Olga Podolich,^d Bertram Brenig,^e Sandeep Tiwari,^f Vasco Azevedo,^f  Aristóteles Góes-Neto,^g  Oleg Reva,^h Natalia Kozyrovska,^d Jean-Pierre de Vera,ⁱ  Bong-Soo Kim^a

^aDepartment of Life Science, Multidisciplinary Genome Institute, Hallym University, Chuncheon, Republic of Korea

^bInstitute for Liver and Digestive Diseases, Hallym University, Chuncheon, Republic of Korea

^cInstitute of Integrative Omics and Applied Biotechnology (IIOAB), Nonakuri, Purba Medinipur, West Bengal, India

^dInstitute of Molecular Biology and Genetics of NASU, Kyiv, Ukraine

^eInstitute of Veterinary Medicine, University of Göttingen, Göttingen, Germany

^fLaboratory of Cellular and Molecular Genetics, Department of General Biology, Institute of Biological Sciences, Federal University of Minas Gerais, Belo Horizonte, Brazil

^gMolecular and Computational Biology of Fungi Laboratory, Department of Microbiology, Institute of Biological Sciences, Federal University of Minas Gerais, Belo Horizonte, Brazil

^hCentre for Bioinformatics and Computational Biology, Department of Biochemistry, Genetics and Microbiology, University of Pretoria, Pretoria, South Africa

ⁱMicrogravity User Support Center (MUSC), German Aerospace Center (DLR), Cologne, Germany

ABSTRACT Kombucha is a traditional tea fermented by symbiotic microbiota, and it has been known as a functional fermented product. Here, we report four microbial metagenome-assembled genome sequences (MAGs) reconstructed from the microbiomes in kombucha exposed to a Mars-like environment outside the International Space Station.

Kombucha is a traditional sugared tea that is fermented using a symbiotic community of acetic acid bacteria and osmophilic acid-tolerant yeasts. Kombucha microbial community (KMC) samples, composed of bacterial and eukaryotic microorganisms, were used as a part of the BIOMEX (BIology and Mars EXperiment) project to understand the influence of exposure to spaceflight and Mars-like conditions on microbiota (1, 2).

Dried KMC pellicles of ecotype IMBG-1 (from the collection of the Institute of Molecular Biology and Genetics of NASU (IMBG), Kyiv, Ukraine) were exposed to Mars-like conditions simulated on low Earth orbit in the three-layer sample carrier mounted on the EXPOSE-R2 facility outside the International Space Station (ISS) (1). During a 2.5-year exposure at the ISS (18 months outside the station and 7 months inside), samples located on the unprotected top level received UV rays (>200 nm), cosmic ionizing radiation, and other fluencies (2). The middle and bottom levels were protected from UV radiation; however, they were maintained in the Mars-like atmosphere and pressure. During the exposure period, all the KMC samples were kept dried with temperature fluctuations. Analogically prepared KMC samples were maintained under laboratory conditions as references.

The KMC samples exposed to Mars-like conditions and laboratory references were reactivated as previously described (1). The growth rates, cellulose-based pellicle yields, and enzyme activities in the KMC samples were different from those in the ground reference samples after reactivation (3, 4). The KMC specimens that had been exposed to Mars-like UV radiation showed major functional alterations compared to the UV-protected samples. However, the genes sequenced from the dominant strains in the KMCs remained unaltered, meaning that the kombucha samples were ecologically resilient. Therefore, we analyzed the metagenome-assembled genome sequences (MAGs) of the dominant species to understand the effect of space/Mars-like conditions on the microecosystem of the KMCs.

Citation Lee I, Barh D, Podolich O, Brenig B, Tiwari S, Azevedo V, Góes-Neto A, Reva O, Kozyrovska N, de Vera J-P, Kim B-S. 2021. Metagenome-assembled genome sequences obtained from a reactivated kombucha microbial community exposed to a Mars-like environment outside the International Space Station. *Microbiol Resour Announc* 10:e00549-21. <https://doi.org/10.1128/MRA.00549-21>.

Editor David A. Baltrus, University of Arizona

Copyright © 2021 Lee et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Debmalya Barh, dr.barh@gmail.com, or Bong-Soo Kim, bkim79@hallym.ac.kr.

Received 27 May 2021

Accepted 16 August 2021

Published 9 September 2021

TABLE 1 General features of metagenome-assembled genomes from kombucha microbial community

MAG	Taxonomic classification	Genome size (bp)	No. of contigs	No. of genes	<i>N</i> ₅₀ (bp)	GC content (%)	CMP ^a (%)	CNT ^a (%)	Mean coverage (x)	BioSample accession no.	GenBank accession no.
kmcMAG001	<i>Komagataeibacter hansenii</i>	4,214,122	108	3,779	79,588	60.14	94.65	4.98	138.26	SAMN16843767	JAEORC0000000000
kmcMAG002	<i>Komagataeibacter rhaeticus</i>	3,326,596	90	3,040	62,857	63.35	99.11	1.69	26.01	SAMN16843789	JAEORD0000000000
kmcMAG003	<i>Komagataeibacter oboediens</i>	3,370,429	63	3,102	88,723	62.04	98.77	1.49	47.65	SAMN16843790	JAEORE0000000000
kmcMAG004	<i>Pseudomonas</i> sp.	6,200,971	143	5,716	66,800	60.57	97.69	2.41	2.31	SAMN16931596	JAEORF0000000000

^a The completeness (CMP) and contamination (CNT) were determined using CheckM v1.1.3.

For the shotgun metagenomic analyses, we used the returned reactivated KMCs (exposed on the three-level-sample carrier), as well as the initial KMC ecotype, which had been saved for sample preparation in the space experiment, and the laboratory kept the desiccated KMC samples used as references (1, 5). For the metagenomic DNA extraction, the KMC samples were cultured in 50 ml black tea-sugar medium until the stage of biofilm formation (5 days), as described in a previous study (5). The biofilm pellicle was removed, and planktonic cells were precipitated using a centrifuge.

Metagenomic DNA was extracted from the planktonic cells of the KMC samples using the innuSPEED bacteria/fungi DNA isolation kit (Analytik Jena AG, Germany). A sequencing library was prepared using a NEBNext Fast DNA fragmentation and library preparation kit (New England BioLabs, Ipswich, MA, USA). Paired-end sequencing (2×150 bp) was performed using a HiSeq 2500 instrument (Illumina, USA).

The raw sequences were filtered for quality control using SolexaQA (6), and the low-quality sequences and artificially duplicated reads (ARDs) were removed using Illumina-Utils (7) and duplicate read inferred sequencing error estimation (DRISSE) (8), respectively. A total of 497,559,912 reads (74,633,986,800 bp) were assembled using MEGAHIT v1.2.9 (9), resulting in 7,303 metagenomic contigs with a length of 65,241,169 bp. After removal of the short reads ($<1,000$ bp), a total of 1,435 contigs were used for the following binning process. The MAGs were reconstructed using the Anvi-refine program in Anvi'o v6.2 (10). The metagenomic contigs were binned based on the single-copy core gene (SCG) set of Kaiju v1.6.2 (11) in the Anvi'o environment. The completeness and contamination of the MAGs were determined using CheckM v1.1.3 (12). Only the MAGs with a completeness of more than 90% and a contamination rate of less 5% were considered for further analysis. Four bacterial MAGs were obtained after a quality check, and the taxonomic classification was detailed by comparing the average nucleotide identity (ANI) with closely related strains in the NCBI Assembly database using the OrthoANI tool (13). Closely related strains were selected based on the binning results. Default parameters were used for all software unless otherwise specified. Three MAGs were classified as *Komagataeibacter*, and one MAG was classified as *Pseudomonas*. The MAG information is summarized in Table 1. The genome sizes of the three *Komagataeibacter* MAGs ranged from 3,326,596 to 4,214,122 bp, and the GC contents ranged from 60.14% to 63.35%. The genome size of the *Pseudomonas* MAG was 6,200,971 bp, and the GC content was 60.57%. Additional genome characterization and functional annotation were automatically performed using the Prokaryotic Genome Annotation Pipeline (PGAP) (14). These MAGs will provide insights into the effect of Mars-like conditions on the microbes in kombucha.

Data availability. The raw sequence data were deposited at the National Center for Biotechnology Information (NCBI) database under the project numbers [PRJNA636820](#), [PRJNA636837](#), [PRJNA636891](#), [PRJNA637016](#), and [PRJNA637018](#). The MAGs are available at GenBank under the BioSample accession numbers [SAMN16843767](#), [SAMN16843789](#), [SAMN16843790](#), and [SAMN16931596](#) and under the accession numbers [JAEORC000000000](#), [JAEORD000000000](#), [JAEORE000000000](#), and [JAEORF000000000](#), respectively.

ACKNOWLEDGMENTS

This study was supported by the Bio & Medical Technology Development Program of the National Research Foundation of Korea (NRF), funded by the South Korean government (MSIT) (2017M3A9F3043837), and the Korea Health Technology R&D Project through the Korea Health Industry Development Institute (KHIDI), funded by the Ministry of Health & Welfare (HP20C0082). The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

REFERENCES

- Podolich O, Kukharensko O, Haidak A, Zaets I, Zaika L, Storozhuk O, Palchikovskaya L, Orlovskaya I, Reva O, Borisova T, Khirunenko L, Sosnin M, Rabbow E, Kravchenko V, Skoryk M, Kremenskoy M, Demets R, Olsson-Francis K, Kozyrovska N, de Vera J-PP. 2019. Multimicrobial kombucha culture tolerates Mars-like conditions simulated on low-Earth orbit. *Astrobiology* 19:183–196. <https://doi.org/10.1089/ast.2017.1746>.
- de Vera J-P, Alawi M, Backhaus T, Baque M, Billi D, Bottger U, Berger T, Bohmeier M, Cockell C, Demets R, de la Torre Noetzel R, Edwards H, Elsaesser A, Faglierone C, Fiedler A, Foing B, Foucher F, Fritz J, Hanke F, Herzog T, Horneck G, Hubers H-W, Huwe B, Joshi J, Kozyrovska N, Kruchten M, Lasch P, Lee N, Leuko S, Leya T, Lorek A, Martinez-Frias J, Meessen J, Moritz S, Moeller R, Olsson-Francis K, Onofri S, Ott S, Pacelli C, Podolich O, Rabbow E, Reitz G, Rettberg P, Reva O, Rothschild L, Sancho LG, Schulze-Makuch D, Selbmann L, Serrano P, Szwedzyk U, et al. 2019. Limits of life and the habitability of Mars: the ESA space experiment BLO-MEX on the ISS. *Astrobiology* 19:145–157. <https://doi.org/10.1089/ast.2018.1897>.

3. Podolich O, Kukhareno O, Zaets I, Orlovskaya I, Palchykovskaya L, Zaika L, Sysoliatin S, Zubova G, Reva O, Galkin M, Horid'ko T, Kosiakova H, Borisova T, Kravchenko V, Skoryk M, Kremensky M, Ghosh P, Barh D, Goes-Neto A, Azevedo V, de Vera J-P, Kozyrovska N. 2020. Fitness of outer membrane vesicles from *Komagataei-bacter intermedius* is altered under the impact of simulated Mars-like stressors outside the International Space Station. *Front Microbiol* 11:1268. <https://doi.org/10.3389/fmicb.2020.01268>.
4. Orlovskaya I, Podolich O, Kukhareno O, Zaets I, Reva O, Khirunenko L, Zmejkoski D, Rogalsky S, Barh D, Tiwari S, Kumavath R, Goes-Neto A, Azevedo V, Brenig B, Ghosh P, de Vera J-P, Kozyrovska N. 2021. Bacterial cellulose retains robustness but its synthesis declines after exposure to a Mars-like environment simulated outside the International Space Station. *Astrobiology* 21:706–717. <https://doi.org/10.1089/ast.2020.2332>.
5. Goes-Neto A, Kukhareno O, Orlovskaya I, Podolich O, Imchen M, Kumavath R, Kato RB, de Carvalho DS, Tiwari S, Brenig B, Azevedo V, Reva O, de Vera J-PP, Kozyrovska N, Barh D. 2021. Shotgun metagenomic analysis of kombucha mutualistic community exposed to Mars-like environment outside the International Space Station. *Environ Microbiol* 23:3727–3742. <https://doi.org/10.1111/1462-2920.15405>.
6. Cox MP, Peterson DA, Biggs PJ. 2010. SolexaQA: at-a-glance quality assessment of Illumina second-generation sequencing data. *BMC Bioinformatics* 11:485. <https://doi.org/10.1186/1471-2105-11-485>.
7. Eren AM, Vineis JH, Morrison HG, Sogin ML. 2013. A filtering method to generate high quality short reads using Illumina paired-end technology. *PLoS One* 8:e66643. <https://doi.org/10.1371/journal.pone.0066643>.
8. Gomez-Alvarez V, Teal TK, Schmidt TM. 2009. Systematic artifacts in metagenomes from complex microbial communities. *ISME J* 3:1314–1317. <https://doi.org/10.1038/ismej.2009.72>.
9. Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. 2015. MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics* 31:1674–1676. <https://doi.org/10.1093/bioinformatics/btv033>.
10. Eren AM, Esen OC, Quince C, Vineis JH, Morrison HG, Sogin ML, Delmont TO. 2015. Anvi'o: an advanced analysis and visualization platform for 'omics data. *PeerJ* 3:e1319. <https://doi.org/10.7717/peerj.1319>.
11. Menzel P, Ng KL, Krogh A. 2016. Fast and sensitive taxonomic classification for metagenomics with Kaiju. *Nat Commun* 7:11257. <https://doi.org/10.1038/ncomms11257>.
12. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25:1043–1055. <https://doi.org/10.1101/gr.186072.114>.
13. Lee I, Ouk Kim Y, Park S-C, Chun J. 2016. OrthoANI: an improved algorithm and software for calculating average nucleotide identity. *Int J Syst Evol Microbiol* 66:1100–1103. <https://doi.org/10.1099/ijsem.0.000760>.
14. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.